### **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



#### INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:		(11) International Publication Number: WO 00/15220
A61K 31/35	A1	(43) International Publication Date: 23 March 2000 (23.03.00)
(21) International Application Number: PCT/SE  (22) International Filing Date: 13 September 1999 (  (30) Priority Data: 9803158-6 16 September 1998 (16.09.9)  (71) Applicant (for all designated States except US): AKTIEBOLAG [SE/SE]; S-151 85 Södertälje (SE  (72) Inventors; and (75) Inventors/Applicants (for US only): EVENDE [GB/US]; Astra Arcus USA, Inc., P.O. Box Rochester, NY 14603 (US). THORBERG, S [SE/SE]; Astra Arcus AB, S-151 85 Södertälje (SE)  (74) Agent: ASTRA AKTIEBOLAG; Intellectual Property S-151 85 Södertälje (SE).	13.09.9  8) S  ASTR  1).  N, Join 2089  eth-Old  E).	BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW ML, MR, NE, SN, TD, TG).  Published  With international search report.  With amended claims.

(54) Title: A NEW COMPOSITION

#### (57) Abstract

This invention relates to a composition comprising (R)-5-carbamoyl-8- fluoro-3-N,N- dicyclobutylamino-3, 4-dihydro-2II- 1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)- 5-phthlalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, the preparation thereof, pharmaceutical formulations containing said composition, use of and a method of treatment of affective disorders such as mood disorders and anxiety disorders with said composition as well as a kit containing said composition.

### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania .	SK	Slovakia
ΛT	Austria	FR	France	LU	Luxembourg	SN	Scnega!
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	G.N	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ.	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

WO 00/15220 PCT/SE99/01599

#### A NEW COMPOSITION

#### Field of the Invention

The present invention relates to a composition which comprises (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyrans in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof. The present invention also relates to a process for the preparation of the inventive composition, pharmaceutical formulations containing said composition and to the use of said composition either by concomitant administration or by separate administration as an improvement of the treatment of affective disorders such as depression, anxiety, obsessive compulsive disorder (OCD), etc.

15

20

#### **Background of the Invention**

Today, it is generally considered that antidepressants take 2-4 weeks to reach full clinical effect. In contrast, the side effects occur immediately. Thus, slow onset of action of antidepressants leads to a vulnerable period for patients in which they experience the side effects, but not the therapeutic effects of drugs. There is often a heavy burden on the treating physician to persuade the patient to continue with the treatment during this period. Furthermore, in suicidal patients, as the onset of action is gradual, initiative may be regained without the experiencing of full reversal of symptoms, leaving a window of risk for suicide and a frequent requirement for hospitalization. An antidepressant with fast onset of action would not only be beneficial due to the faster symptom reduction, but would also be more acceptable to patients and physicians and reduce the need for and duration of hospitalization. The same long period to reach full clinical effect has been shown in the treatment of other affective disorders such as anxiety and OCD.

#### Prior art

20

In WO 96/33710 is disclosed that the compound (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran which has high affinity to 5-HT receptors-and antagonizes 5-HT<sub>1A</sub> mediated responses induces a rapid improvement of depressed patients treated with serotonin reuptake inhibitors.

#### Summary of the Invention

The present invention is directed to a new composition comprising of the specific 5-HT<sub>1</sub>Aantagonist (R)-5-carbamoyl-8-fluoro-3-N,N-disubstituted-amino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and the specific 5-HT reuptake inhibitor (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof. Said composition attains a faster onset of action and consequently, provides a more efficacious treatment of the patients suffering from affective disorders, particularly depression.

It has been shown in animal studies that acute administration of selective 5-HT reuptake inhibitors (SSRIs) decreases the electrical impulse propagation in 5-HT neurones via a negative feedback reaction probably mediated by collateral 5-HT axons releasing 5-HT in raphé nuclei. By inhibiting the somatodendritic 5-HT<sub>1A</sub> autoreceptors in the raphé nuclei the selective antagonists counteract the decrease in propagetion caused by 5-HT reuptake inhibitors. This indicates that a selective blockade of somatodendritic autoreceptor i.e. 5-HT<sub>1A</sub> antagonist may have a clinical potential to improve the efficacy of 5-HT reuptake inhibitors (SSRIs) and offer a new rationale for rapid onset of effect in the treatment of affective disorders, for instance the antidepressant actions.

25

30

The compound (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2*H*-1-benzopyran in the form of the free base, and pharmaceutically acceptable salts thereof as disclosed herein is described in WO 95/11891, as a selective 5-HT1A receptor antagonist.

The enantiomer (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5phthalancarbonitrile disclosed herein, and which is stated to be a 5-HT reuptake inhibitor is described in US 4,943,590.

The (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran is in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof. Both organic and inorganic acids can be employed to form nontoxic pharmaceutically acceptable acid addition salts of the compounds of this invention. Illustrative acids are sulfuric, nitric, phosphoric, oxalic, hydrochloric, formic, hydrobromic, citric, acetic, lactic, tartaric, dibenzoyltartaric, diacetyltartaric, pamoic, ethanedisulfonic, sulfamic, succinic, propionic, glycollic, malic, gluconic, pyruvic, phenylacetic, 4-aminobenzoic, anthranilic, salicylic, 4-aminosalicylic, 4-hydroxybenzoic, 3,4-dihydroxybenzoic, 3,5-dihydroxybenzoic, 3-hydroxy-2-naphthoic, nicotinic, methanesulfonic, ethanesulfonic, hydroxyethanesulfonic, benzenesulfonic, p-toluenesulfonic, sulfanilic, naphthalenesulfonic, ascorbinic, cyclohexylsulfamic, fumaric, maleic and benzoic acids. These salts are readily prepared by methods known in the art.

The (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran possesses a high affinity to the specific subgroup of 5-HT<sub>1A</sub> receptor in the CNS and acts as an antagonist on that 5-HT<sub>1A</sub> receptor, and as also shows favourable bioavailability after oral administration.

The composition according to the present invention may exist in one pharmaceutical formulation comprising of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-

phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof. Alternatively, the composition may exist in two different pharmaceutical formulations, one for (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable-salt and/or solvate thereof and one for (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof. The pharmaceutical formulation may be in the form of tablets or capsules, powders, mixtures, solutions or other suitable pharmaceutical formulation forms such as patches and nasal formulations.

10

15

20

5

such

The composition of the present invention can be prepared by that (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is incorporated into the same pharmaceutical formulation as (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof by e.g. mixing in a conventional way.

The present invention also includes a method of improving the onset of therapeutic action by concomitant administration of a composition comprising (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof and (+)-1-[3-(dimethylamino)-propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.

A further embodiment of the present invention is a kit containing a dosage unit of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclbutylamino-3,4-dihydro-2H-1-benzopyrans in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and a dosage unit of (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, optionally with instructions for use.

#### Pharmaceutical formulations

20

According to the present invention the compounds in the composition will normally be administered orally, rectally, transdermally, nasally or by injection, in the form of pharmaceutical formulations comprising the active ingredient either as a free base, a solvate e.g. a hydrate or a pharmaceutically acceptable non-toxic acid addition salt, e.g. the hydrochloride, hydrobromide, lactate, acetate, phosphate, sulfate, sulfamate, citrate, tartrate, oxalate and the like in a pharmaceutically acceptable dosage form. The dosage form may be a solid, semisolid or liquid formulation. Usually the active substances will constitute between 0.1 and 99% by weight of the formulation, more specifically between 0.5 and 20% by weight for formulations intended for injection and between 0.2 and 50% by weight for formulations suitable for oral administration.

The pharmaceutical formulation comprises the active ingredients, optionally in association with adjuvants, diluents, excipients and/or inert carriers.

To produce pharmaceutical formulations of the composition of the invention in the form of dosage units for oral application, the selected compounds may be mixed with a solid excipient, e.g. lactose, saccharose, sorbitol, mannitol, starches such as potato starch, corn starch or amylopectin, cellulose derivatives, a binder such as gelatin or polyvinylpyrrolidone, disintegrants e.g. sodium starch glycolate, cross-linked PVP and cross-caramellose sodium; and a lubricant such as magnesium stearate, calcium stearate, polyethylene glycol, waxes, paraffin, and the like, and an antisticking agent such as talc or colloidal silicon dioxide, and then compressed into tablets. If coated tablets are required, the cores, prepared as described above, may be coated with a polymer known to the man skilled in the art e.g. HPMC, HC or other cellulose derivatives or PVP, wherein the polymer is dissolved in water or a readily volatile organic solvent or mixture of organic solvents. Alternatively, the tablets can be coated with a concentrated sugar solution which may contain e.g. gum arabic, gelatine, talcum, titanium dioxide, and the like. Dyestuffs

may be added to these coatings for instance in order to readily distinguish between tablets containing different active substances or different amounts of the active compounds.

For the formulation of soft gelatin capsules, the active substances may be admixed with e.g. a vegetable oil or polyethylene glycol. Hard gelatine capsules may contain granules of the active substances using any of the above mentioned excipients for tablets e.g. lactose, saccharose, sorbitol, mannitol, starches (e.g. potato starch, corn starch or amylopectin), cellulose derivatives, plasticizers, polyetheneglycole, waxes, lipids or gelatin. Also liquids or semisolids of the drug can be filled into hard gelatin capsules.

10

15

20

25

30

Dosage units for rectal application can be solutions or suspensions or can be prepared in the form of suppositories comprising the active substances in a mixture with a neutral fatty base, or gelatin rectal capsules comprising the active substances in admixture with vegetable oil or paraffin oil. Liquid formulations for oral application may be in the form of solutions, syrups or suspensions, for example solutions containing from about 0.2% to about 20% by weight of the active substances herein described, the balance being sugar and a mixture of ethanol, water, glycerol and propylene glycol. Optionally such liquid formulations may contain colouring agents, flavouring agents, saccharin and carboxymethyl-cellulose as a thickening agent or other excipients known to a person skilled in the art.

Solutions for parenteral applications by injection can be prepared in an aqueous solution of a water-soluble pharmaceutically acceptable salt of the active substances, preferably in a concentration of from about 0.5% to about 10% by weight. These solutions may also contain stabilizing agents and/or buffering agents and may conveniently be provided in various dosage unit ampoules.

Suitable daily doses of the active compounds in the composition of the invention in therapeutic treatment of humans are about 0.01-100 mg/kg bodyweight for peroral administration and 0.001-100 mg/kg bodyweight for parenteral administration. The daily

5

doses of the active ingredient (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof may very well differ from the daily doses of the active ingredient (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof but the doses can also be the same for both of the active ingredients.

#### Medical and Pharmaceutical Use

In a further aspect the present invention provides the use of the composition of (R)-5-10 carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof and (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and the use in the treatment of 5-hydroxytryptamine mediated disorders, such as affective disorders. Examples of affective disorders are disorders in the CNS such as mood disorders (depression, major depressive episodes, dysthymia, seasonal affective disorder, depressive phases of bipolar disorder), anxiety disorders (obsessive compulsive disorder, panic disorder with/without agoraphobia, social phobia, specific phobia, generalized anxiety disorder, posttraumatic stress disorder), personality disorders (disorders of impulse control, 20 trichotellomania) and sleep disorders. Other disorders in the CNS such as eating disorders(obesity, anorexia, bulimia), premenstrual syndrome, sexual disturbances, alcoholism, tobacco abuse, autism, attention deficit, hyperactivity disorder, migraine, memory disorders (age associated memory impairment, presenile and senile dementia such as Alzheimer's disease), pathological aggression, schizophrenia, endocrine disorders (e g hyperprolactinaemia), stroke, dyskinesia, Parkinson's disease, thermoregulatory disorders, pain and hypertension may also be treated with the combination described herein. Examples of other hydroxytryptamine mediated disorders are urinary incontinence, vasospasm and growth control of tumors (e g lung carcinoma) and it may be possible to treat those with the combination described herein as well.

#### Pharmacology

Potentiation of the 5 HT<sub>1A</sub> autoreceptor blocking effekt of 5-HT of (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phtalancarbonitrile by using of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran.

#### Materials and methods

#### Animals

The studies were carried out in male Sprague-Dawley rats (290-450g; B&K Universal, Sollentuna, Sweden). The animals were housed for at least 3 weeks after arrival until used in the experiments.

#### Methods

The studies were carried out by means of intra-cerebral microdialysis in awake rats. To assess any putative regional differences between dorsal and median rapheinnervated 5-HT projection areas, dialysis probes were simultaneously implanted both into the frontal cortex (FCx) and dorsal hippocampus (DH).

### 20 Microdialysts

25

The rats were anaesthetised with a mixture of ketamine HCl (67 mg/kg intraperitoneal (IP); Ketalar Park-Davis) and xylazine HCl (13 mg/kg IP; Rompun Bayer-Leverkusen). Ushaped microdialysis probes (total dialysis fibre length 4 mm, OD 220µm) were stereotaxically implanted in the frontal cortex (FCx) and dorsal hippocampus (DH); probe tips at AP +3.5, ML -3.0, DV -4.2 and -4.3, ML +2.5, DV -4.2, respectively, vs. bregma and dura surface (Paxinos, et al, in The Rat Brain in Stereotaxic Coordinates, 2nd Ed., Academic Press, San Diego (1996)). The microdialysis studies were performed in conscious animals after a 40-48 h recovery period, during which they were kept individually. Food and water were allowed ad libitum in the plastic cages subsequently used in the experimental sessions. On the day of the experiment, the probe inlets were

connected to a syringe perfusion pump (CMA/100; CMA Microdialysis AB, Sweden), delivering artificial CSF (Hjorth, S., J. Neurochem. 60:776-779 (1993)) at a speed of 1.3µl/min. Twenty-min dialysate fractions were collected from the probe outlet tubing, and immediately analysed for 5-HT and 5-HIAA by standard HPLC-EC methods. After the perfusion was commenced, a period of 2-3 h was allowed to establish stable baseline dialysate levels of 5-HT, prior to drug treatment(s).

#### **CLAIMS**

20

- 1. A composition comprising of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.
- 2. Use of the composition according to claim 1 for the manufacture of a medicament for the treatment of 5-HT mediated disorders.
  - 3. The use according to claim 2 for the manufacture of a medicament for the treatment of affective disorders.
- 4. The use according to claim 3 for the manufacture of a medicament for the treatment of mood disorders.
  - 5. The use according to claim 4 for the manufacture of a medicament for the treatment of depression.
  - 6. A method for the treatment of 5-HT mediated disorders by administering to a patient suffering therefrom the composition defined in claim 1.
  - 7. The method according to claim 6 for the treatment of affective disorders.
  - 8. The method according to claim 7 for the treatment of mood disorders.
  - 9. The method according to claim 8 for the treatment of depression.

WO 00/15220 PCT/SE99/01599

11

- 10. A method of improving the onset of therapeutic action by concomitant administration of a composition defined in claim 1.
- 11. A pharmaceutical formulation wherein the active ingredients are those in the composition defined in claim 1, optionally in association with adjuvants, excipients and/or inert carriers.
- 12. A pharmaceutical formulation according to claim 11 wherein (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is concomitantly administered with (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.
- 13. A pharmaceutical formulation according to any one of claims 11-12 for use in the treatment of 5-HT mediated disorders.
  - 14. A pharmaceutical formulation according to any one of claims 11-12 for use in the treatment of affective disorders.
- 15. A pharmaceutical formulation according to any one of claims 11-12 for use in the treatment of mood disorders.

- 16. A pharmaceutical formulation according to any one of claims 11-12 for use in the treatment of depression.
- 17. A process for the preparation of the composition according to claim 1 whereby (R)-5-carbamoyl-8-fluoro-3-N,N-dicycl0butylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is incorporated into the same pharmaceutical formulation as (+)-1-[3-(dimethylamino)propyl]-1-(p-

fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.

18. A process for the preparation of the composition according to claim 1 whereby (R)-5- carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is in a one pharmaceutical formulation is combined with (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is in a different pharmaceutical formulation.

10

15

5

19. A kit containing a dosage unit of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and a dosage unit of (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, optionally with instructions for use.

#### **AMENDED CLAIMS**

[received by the International Bureau on 15 February 2000 (15.02.00); original claims 1-19 replaced by amended claims 1-22 (3 pages)]

- 1. A composition comprising of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.
- 2. Use of the composition according to claim 1 for the manufacture of a medicament for the treatment of 5-HT mediated disorders.
  - 3. The use according to claim 2 for the manufacture of a medicament for the treatment of affective disorders.
- 4. The use according to claim 3 for the manufacture of a medicament for the treatment of mood disorders.
  - 5. The use according to claim 4 for the manufacture of a medicament for the treatment of depression.
  - 6. The use according to claim 2 in the manufacture of a medicament in the prevention or in the treatment of urinary incontinence.
  - 7. A method for the treatment of 5-HT mediated disorders by administering to a patient suffering therefrom the composition defined in claim 1.
    - 8. The method according to claim 7 for the treatment of affective disorders.
    - 9. The method according to claim 8 for the treatment of mood disorders.

20

5

- 10. The method according to claim 9 for the treatment of depression.
- 11. A method according to claim 7 for the prevention or the treatment of urinary incontinence.
- 12. A method of improving the onset of therapeutic action by concomitant administration of a composition defined in claim 1.
- 13. A pharmaceutical formulation wherein the active ingredients are those in the composition defined in claim 1, optionally in association with adjuvants, excipients and/or inert carriers.
  - 14. A pharmaceutical formulation according to claim 13 wherein (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is concomitantly administered with (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.
- 15. A pharmaceutical formulation according to any one of claims 13-14 for use in the treatment of 5-HT mediated disorders.
  - 16. A pharmaceutical formulation according to any one of claims 13-14 for use in the treatment of affective disorders.
  - 17. A pharmaceutical formulation according to any one of claims 13-14 for use in the treatment of mood disorders.
  - 18. A pharmaceutical formulation according to any one of claims 13-14 for use in the treatment of depression.

- 19. A pharmaceutical formulation according to any one of claims 13-14 for use in the treatment of urinary incontinence.
- 20. A process for the preparation of the composition according to claim 1 whereby (R)-5-carbamoyl-8-fluoro-3-N,N-dicycl0butylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is incorporated into the same pharmaceutical formulation as (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof.
  - 21. A process for the preparation of the composition according to claim 1 whereby (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is in a one pharmaceutical formulation is combined with (+)-1-[3-(dimethylamino)propyl]-1-(p-fluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof is in a different pharmaceutical formulation.
- 22. A kit containing a dosage unit of (R)-5-carbamoyl-8-fluoro-3-N,N-dicyclobutylamino-3,4-dihydro-2H-1-benzopyran in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, and a dosage unit of (+)-1-[3-(dimethylamino)propyl]-1-(pfluorophenyl)-5-phthalancarbonitrile in the form of the free base, or a pharmaceutically acceptable salt and/or solvate thereof, optionally with instructions for use.

### INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 99/01599

A. CLASS	SIFICATION OF SUBJECT MATTER		
IPC7: A	61K 31/35  Distributional Patent Classification (IPC) or to both nat	ional classification and IPC	·
	S SEARCHED		
	ocumentation searched (classification system followed by	classification symbols)	·
IPC7: A		to the state of th	she Golda asseshed
	ion searched other than minimum documentation to the	extent that such documents are included if	i die fields sealched
	I,NO classes as above		
Electronic da	ata base consulted during the international search (name	of data base and, where practicable, search	terms used)
C. DOCU	MENTS CONSIDERED TO BE RELEVANT		·
Category*	Citation of document, with indication, where app	ropriate, of the relevant passages	Relevant to claim No.
A	WO 9633710 A1 (ASTRA AKTIEBOLAG) (31.10.96)	, 31 October 1996	1-5,11-19
L			
			ĺ
			! 
			i
			1
	L. C.P.	C Suppose family again	_
Furth	ter documents are listed in the continuation of Box		
"A" docum	eategones of cited documents cat defining the general state of the art which is not considered of particular relevance	"I" later document published after the int date and not in conflict with the appl the principle or theory underlying the	ication but cited to understand
"E" erlier o	focument but published on or after the international filing date ent which may throw doubts on priority claim(s) or which is	"X" document of particular relevance: the considered nevel or cannot be considered when the document is taken along	ered to involve an inventive
ested to special	o establish the publication date of another diation or other reason (as specified)	"Y" document of particular relevance: the	claimed invention cannot be
meany		considered to involve an inventive ste combined with one or more other ste being obvious to a person skilled in t	h documents, such commonation
	ent published prior to the international filing date but fater than only date claimed	'&' document member of the same paten	
Date of th	e actual completion of the international search	Date of mailing of the international	
1 20 0	anhan 1000	2 2	2 -01- 2000
Name and	ember 1999 I mailing address of the ISA	Authorized officer	
Swedish	Patent Office	0" == 1/21 = - / 1/0	
1	5, S-102 42 STOCKHOLM No. +46 8 666 02 86	Göran Karlsson / MR Telephone No. +46 8 782 25 00	

Form PCT ISA 210 (second sheet) (July 1992)

### INTERNATIONAL SEARCH REPORT

Internation... application No. PCT/SE99/01599

Box l	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This inte	rnational search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:
1. 🔯	Claims Nos.: 6-10 because they relate to subject matter not required to be searched by this Authority, namely:  A method for treatment of the human or animal body by therapy, see Rule 39.1
2.	Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  .
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This Inte	mational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.	As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	on Protest  The additional search fees were accompanied by the applicant's protest.  No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

			02/12/99	PC1/SE	99/01599
Patent document cited in search report	Publication date		Patent family member(s)		Publication date
0 9633710 A1	31/10/96	AU	696356	В	10/09/98
		AU	5520296	A	18/11/96
•		BR	9608252	A	04/05/99
		CA	2218181	A	31/10/96
		CZ	9703356	A	17/06/98
		EP	0807241		19/11/97
		EP	0822813	A	11/02/98
		ΙL	117988	D	00/00/00
		JP	11504037	T	06/04/99
		NO	973445	A	25/07/97
		NO	974921		18/12/97
		NZ	306602		29/04/99 ·
		PL	323083	A	02/03/98
		SE	9501567	D	00/00/00
		SK	142997	A	02/12/98
		US -	5962514	A	05/10/99
		ZA	9602982	Δ	28/10/96

Form PCT ISA 210 (patent family annex) (July 1992)